

WORKSHEET 3 : Differentiation I

1. Find the points where the following functions have horizontal tangent.

a) $f(x) = x^3 + 1$ b) $f(x) = 1/x^2$ c) $f(x) = x + \sin x$
d) $f(x) = \sqrt{x-1}$ e) $f(x) = e^x - x$ f) $f(x) = \sin x + \cos x$

a) $x=0$; b) never; c) $x = \pi + 2k\pi$; d) never; e) $x = 0$; f) $x = \frac{\pi}{4} + k\pi$.

2. (*) Prove that the tangent lines to the graphs of $y = x$ and $y = 1/x$ in their points of intersection are perpendicular to each other.

3. In what point is the tangent to the curve $y^2 = 3x$ parallel to the line $y = 2x$?

The point of the curve is $(\frac{3}{16}, \frac{3}{4})$.

4. (*) Calculate the intersection point with the x axis of the tangent line to the graph of $f(x) = x^2$ in the point $(1, 1)$.

The intersection point is $x = 1/2$.

5. Calculate a so that the tangent to the graph of $f(x) = a/x + 1$ in the point $(1, f(1))$ intersects the horizontal axis in $x = 3$.

Therefore, the intersection point will be $x=3$ when $a=1$.

6. (*) Find the tangent and normal lines to $f(x) = \arctan\left(\frac{\sin x}{1+\cos x}\right)$ in $x = 0$.

Equation of the tangent line: $y - 0 = \frac{1}{2}(x - 0)$; equation of the normal line: $y - 0 = -2(x - 0)$.

7. Find the derivatives of the following functions.

a) $f(x) = (\sin x + \tan 3x)\sin 2x$ b) $f(x) = \frac{x\sqrt{x^2-1}}{2x+6}$
c) $f(x) = 4x^{3/2} \cos 2x$ d) $f(x) = 5x \ln(8x + \sin 2x) + e^{\tan 5x}$

8. (*) Let $f(x) = 2[\ln(1 + g^2(x))]^2$. Using that $g(1) = g'(1) = -1$, calculate $f'(1)$.

$f'(1) = 4 \ln(2)$.

9. (*) Using that $a^b = e^{b \ln a}$, differentiate $f(x) = x^{\sin x}$ and $g(x) = (\sqrt{x})^x$.

$f'(x) = x^{\sin x} (\cos x \cdot \ln x + \sin x/x)$.

$g'(x) = (\sqrt{x})^x (\ln x + 1)/2$.

10. (*) Let $f(x) = \ln(1 + x^2)$ and $g(x) = e^{2x} + e^{3x}$. Calculate $h(x) = f(g(x))$, $v(x) = g(f(x))$, $h'(0)$ and $v'(0)$.

$h(x) = \ln(1 + e^{4x} + e^{6x} + 2e^{5x})$, $h'(0) = 4$

$v(x) = (1 + x^2)^2 + (1 + x^2)^3$, $v'(0) = 0$.

11. Let $f : [-2, 2] \rightarrow [-2, 2]$ be continuous and bijective.

a) Suppose that $f(0) = 0$ and $f'(0) = \alpha$, $\alpha \neq 0$. Find $(f^{-1})'(0)$.

b) Now suppose that $f(0) = 1$ and $f'(0) = \alpha$, $\alpha \neq 0$. Find $(f^{-1})'(1)$.

c) Now suppose that $f(1) = 0$ and $f'(1) = \alpha$, $\alpha \neq 0$. Find $(f^{-1})'(0)$.

a) $(f^{-1})'(0) = \frac{1}{\alpha}$.

b) $(f^{-1})'(1) = \frac{1}{\alpha}$.

c) $(f^{-1})'(0) = \frac{1}{\alpha}$.

12. (*)Supposing that the following equations define y as a differentiable function of x , calculate y' in the given points:

a) $x^3 + y^3 = 2xy$ in $(1, 1)$.

b) $\sin x = x(1 + \tan y)$ in $(\pi, 3\pi/4)$.

c) $x^2 + y^2 = 25$ in $(3, 4), (0, 5)$ and $(5, 0)$.

a) $y' = -1$. b) $y' = \frac{-1}{2\pi}$. c) $y' = \frac{-3}{4}$ in $(3, 4)$. $y' = 0$ in $(0, 5)$. It doesn't exist derivative in $(5, 0)$.

13. Calculate the derivative of the following functions showing where they are not differentiable.

a) (*) $f(x) = \begin{cases} x^2 - 1 & \text{if } x \leq 0 \\ 0 & \text{if } x > 0 \end{cases}$ b) (*) $g(x) = \begin{cases} 1/|x| & \text{if } x \leq -2 \\ (x+2)^2 & \text{if } -2 < x \leq 0 \\ 3 + \sin(x + \frac{\pi}{2}) & \text{if } x > 0 \end{cases}$

c) $h(x) = \begin{cases} \arctan^2 x & \text{if } x \leq 0 \\ \sin^3 x & \text{if } 0 < x \leq 2\pi \\ \sin x & \text{if } 2\pi < x \end{cases}$

a) $f'(x) = \begin{cases} 2x & \text{if } x < 0 \\ 0 & \text{if } x > 0 \end{cases}$ Obviously, f is not differentiable in 0 .

b) $g'(x) = \begin{cases} 1/x^2 & \text{if } x < -2 \\ 2(x+2) & \text{if } -2 < x < 0 \\ \cos(x + \frac{\pi}{2}) & \text{if } x > 0 \end{cases}$ Obviously, f is not differentiable in -2 nor in 0 .

c) $h'(x) = \begin{cases} \frac{2 \arctan x}{1+x^2} & \text{if } x \leq 0 \\ 3 \sin^2 x \cdot \cos x & \text{if } 0 \leq x < 2\pi \\ \cos x & \text{if } 2\pi < x \end{cases}$ Obviously, f is not differentiable in 2π .

14. (*)Find a and b so that the function $f(x) = \begin{cases} 3x + 2 & \text{if } x \geq 1 \\ ax^2 + bx - 1 & \text{if } x < 1 \end{cases}$ is differentiable.

f differentiable in 1 when $a = -3, b = 9$.

15. Apply the mean value theorem to f in the given interval and find the c values of the thesis of the theorem.

a) $f(x) = x^2$ in $[-2, 1]$

b) $f(x) = -2\sin x$ in $[-\pi, \pi]$

c) $f(x) = x^{2/3}$ in $[0, 1]$

d) $f(x) = 2\sin x + \sin 2x$ in $[0, \pi]$

16. Let $f : [a, b] \rightarrow [a, b]$ be a continuous function in $[a, b]$ and differentiable in (a, b) . Prove that, if $f'(x) \neq 1$ in (a, b) , then f has a unique fixed point in $[a, b]$.

17. Prove that the function f has a unique fixed point.

a) $f(x) = 2x + \frac{1}{2}\sin x$

b) $f(x) = 2x + \frac{1}{2}\cos x$

18. (*)Let $f(x) = x^3 - 3x + 3$, $f : [-3, 2] \rightarrow \mathbb{R}$. Determine the global extrema.

The minimum is reached in -3 and the maximum is reached in -1 and in 2 .

19. Let $f : [-5, 5] \rightarrow \mathbb{R}$ such that f reaches the maximum in $x = 2$ and the minimum in $x = -3$. Let $g(x) = -f(-x)$. What can be said about the maximum and the minimum of g ?