

WORKSHEET 5: Integration

1. (*) Calculate the following integrals:

$$\begin{array}{lll}
 \text{a) } \int \frac{x^2 + x + 1}{x\sqrt{x}} dx & \text{b) } \int xe^{-2x} dx & \text{c) } \int \sin^{14} x \cos x dx \\
 \text{d) } \int (x+1)(2-x)^{1/3} dx & \text{e) } \int \frac{x^4}{1+x^5} dx & \text{f) } \int (1 + \frac{1}{x})^3 \frac{1}{x^2} dx \\
 \text{g) } \int \sin^3 x dx & \text{h) } \int xe^{ax^2} dx & \text{i) } \int \frac{1}{3+x^2} dx \\
 \text{j) } \int \frac{\sqrt{x-1}}{1+\sqrt[3]{x-1}} dx & \text{k) } \int \frac{x}{\sqrt{16-x^2}} dx & \text{l) } \int x^4 \ln x dx \\
 \text{m) } \int \frac{dx}{\sqrt[4]{x^3} - \sqrt{x}} & \text{n) } \int (\ln x)^2 dx & \tilde{\text{n) } } \int \frac{40x}{(x-1)^{40}} dx \\
 \text{o) } \int \frac{4x+6}{(x^2+3x+7)^3} dx & \text{p) } \int \frac{2x-6}{(x-2)^2} dx & \text{q) } \int \frac{x^2+1}{x^3-4x^2+4x} dx \\
 \text{r) } \int \frac{2x+1}{x^3+6x} dx & \text{s) } \int \frac{1}{\frac{x^2}{2} - 2x + 4} dx & \text{t) } \int \frac{x^4}{x^4-1} dx
 \end{array}$$

Solution:

$$\begin{array}{l}
 \text{a) } \frac{2}{3}x^{3/2} + 2x^{1/2} - 2x^{-1/2} + C \\
 \text{b) } xe^{-2x}/2 - \frac{1}{4}e^{-2x} + C \\
 \text{c) } (\sin^{15}x)/15 + C \\
 \text{d) } (-3)\frac{3}{4}t^{4/3} + \frac{3}{7}t^{7/3} + C \\
 \text{e) } \frac{1}{5} \ln(1+x^5) + C \\
 \text{f) } (1 + \frac{1}{x})^4/4 + C \\
 \text{g) } -\cos x + (\cos^3 x)/3 + C \\
 \text{h) } \frac{1}{2a} e^{ax^2} + C \\
 \text{i) } \frac{\sqrt{3}}{3} \arctan(\frac{x\sqrt{3}}{3}) + C \\
 \text{j) } 6[\frac{1}{7}\sqrt[6]{(x-1)^7} - \frac{1}{5}\sqrt[6]{(x-1)^5} + \frac{1}{3}\sqrt[6]{(x-1)^3} - \sqrt[6]{(x-1)} + \arctan \sqrt[6]{x-1}] + C \\
 \text{k) } -\sqrt{16-x^2} + C \\
 \text{l) } \frac{1}{5}x^5 \ln x - \frac{1}{25}x^5 + C \\
 \text{m) } 4\sqrt[4]{x} + 4 \ln(\sqrt[4]{x} - 1) + C \\
 \text{n) } x(\ln x)^2 - 2x \ln x + 2x + C \\
 \tilde{\text{n) } } 40[(\frac{-1}{38})(x-1)^{-38} + (\frac{-1}{39})(x-1)^{-39}] + C \\
 \text{o) } 2(x^2+3x+7)^{-2} /(-2) + C \\
 \text{p) } 2 \ln(x-2) + \frac{2}{x-2} + C \\
 \text{q) } \frac{1}{4} \ln x + \frac{3}{4} \ln(x-2) - \frac{5}{2}(x-2)^{-1} + C \\
 \text{r) } \frac{-1}{12} \ln(x^2+6) + 2\frac{\sqrt{6}}{6} \arctan(\frac{x\sqrt{6}}{6}) + \frac{1}{6} \ln x + C \\
 \text{s) } \arctan(\frac{x-2}{2}) + C \\
 \text{t) } x + \frac{1}{4} \ln(x-1) - \frac{1}{4} \ln(x+1) - \frac{1}{2} \arctan(x) + C
 \end{array}$$

2. How many different intersection points can two different primitives of the same function have?

Solution: none.

3. (*) Let $f : [0, 2] \rightarrow \mathbb{R}$ be continuous, increasing in $(0, 1)$, decreasing in $(1, 2)$ and, also, satisfying that: $f(0) = 3$, $f(1) = 5$ and $f(2) = 4$. Between which values can we guarantee that $\int_0^2 f(x) dx$ is located?

Solution: $7 \leq \int_0^2 f(x) dx \leq 10$.

4. (*) Certain company has determined that its marginal cost is $\frac{dC}{dx} = 4(1 + 12x)^{-1/3}$.

Find the cost function if $C = 100$ when $x = 13$.

Solution: $\frac{1}{3}(1 + 12x)^{2/3} \cdot \frac{3}{2} + 100 - \frac{1}{2}(157)^{2/3}$

5. (*) Given that the marginal cost of producing x units is $x + 5$ and the average cost has a minimum in $x = 4$, find the fixed costs of the firm.

Solution: $C_f = 8$.

6. (*) Calculate $F'(x)$ in the following cases:

$$a) \int_x^{x^3} t \cos t dt \quad b) \int_1^{x^2} \sqrt{t^4 + 2t} dt \quad c) \int_1^{x^2} (t^2 - 2t + 5) dt$$

Solution:

$$a) F'(x) = 3x^5 \cos x^3 - x \cos x$$

$$b) F'(x) = \sqrt{x^8 + 2x^2} \cdot 2x$$

$$c) F'(x) = 2x^5 - 4x^3 + 10x$$

7. Calculate $F'(x)$ in the following cases:

$$(a) \int_{-x}^{x^2} \tan^2 t dt, \text{ supposing that } x^2 < \frac{\pi}{2}.$$

$$(b) \int_{x^2}^{2x} f^2(2t) dt, \text{ supposing that } f \text{ is continuous.}$$

Solution:

$$(a) 2x \tan^2 x^2 + \tan^2 x$$

$$(b) 2 f^2(4x) - 2x f^2(2x^2)$$

8. (*) What are the values of x where $F(x) = \int_{-3}^x \frac{t^2 - 4}{3t^2 + 1} dt$ has a local maximum or minimum?

Solution: F reaches a local minimum in 2 and a local maximum in -2.

9. Let $F(x) = \int_{x^2}^{2x} f(t^2) dt$ be such that $f(1) = 1$, $f(2) = f(4) = 4$ and f is continuous. Calculate $F'(1)$.

Solution: $F'(1) = 6$

10. (*) Calculate observing the symmetry of the functions:

$$a) \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^{27} x \cos^{28} x dx \quad b) \int_{-\frac{\pi}{3}}^{\frac{\pi}{3}} (\sqrt[3]{x^5 \cos 3x} + \cos \frac{x}{3} + \tan^3 x) dx$$

Solution:

(a) 0

(b) $6 \sin \frac{\pi}{9}$.

11. Let f be a function with period T , such that $\int_0^T f = b$. Find $\int_a^{a+nT} f$.

Solution: $\int_a^{a+nT} f = \int_0^{0+nT} f = nb$

12. (*) Find the area located between the following curves:

a) $f(x) = x^2 - 4x + 3$, $g(x) = -x^2 + 2x + 3$

b) $f(x) = (x - 1)^3$, $g(x) = x - 1$

c) $f(x) = x^4 - 2x^2 + 1$, $g(x) = 1 - x^2$

Solution:

(a) 9.

(b) $\frac{1}{2}$.

(c) $\frac{4}{15}$.

13. (*) Graph the functions $y = 2e^{2x}$ and $y = 2e^{-2x}$. Calculate the area located between those graphs and the lines $x = -1$ and $x = 1$.

Solution: $2(e^2 + e^{-2} - 2)$

14. Let $f : [1, 3] \rightarrow [2, 4]$ be increasing, continuous and bijective such that $\int_1^3 f dx = 5$. Calculate $\int_2^4 f^{-1}(x) dx$

Solution: 5

15. (a) Given $f : [0, 4] \rightarrow \mathbb{R}$, convex and increasing with values $f(0) = 0$, $f(2) = \alpha$, $f'(2) = \beta$, $f(4) = 16$. Estimate as a function of α and β , the value of $\int_0^4 f(x) dx$.

(b) Given $f : [0, 4] \rightarrow \mathbb{R}$, concave and increasing with values $f(0) = 0$, $f(2) = \alpha$, $f'(2) = \beta$, $f(4) = 2$. Estimate as a function of α and β , the value of $\int_0^4 f(x) dx$.

Solution:

(a) $2(\alpha + \beta) < \int_0^4 f(x) dx < 2\alpha + 32$.

(b) $2\alpha < \int_0^4 f(x) dx < 2(\alpha - \beta) + 4$.

16. The sales of a product are given by the formula $S(t) = 10 + 5 \sin(\frac{\pi t}{6})$ where S is measured in thousands of units and time t in months. Calculate the average sales during the year ($0 \leq t \leq 12$).

Solution: 10

17. Calculate:

$$\begin{array}{lll} a) \int_0^1 \frac{1}{\sqrt{x}} dx & b) \int_0^3 \frac{1}{x^3} dx & c) \int_1^\infty \frac{1}{x^2} dx \\ d) \int_1^\infty e^{-x} dx & e) \int_{-\infty}^\infty \frac{dx}{1+x^2} & f) \int_{-2}^4 \frac{dx}{x^2} \end{array}$$

Solution:

a) 2

b) ∞

c) 1

d) 1

e) π

f) ∞

18. Calculate $\int_0^\infty \frac{dx}{\sqrt{x}(1+x)}$

Solution: π