

1

(a) (10 puntos) Sabiendo que

$$\begin{vmatrix} 1 & 0 & 1 & a \\ 2 & -1 & 1 & b \\ 3 & 4 & 0 & c \\ 1 & -1 & 0 & d \end{vmatrix} = 10,$$

encontrar de forma justificada el valor del determinante

$$\begin{vmatrix} 1 & 0 & 2a+6 & 1 \\ 2 & -1 & 2b & 1 \\ 3 & 4 & 2c & 0 \\ 1 & -1 & 2d & 0 \end{vmatrix}.$$

(b) (10 puntos) Encontrar una de la matrices escalonadas equivalente a la matriz de orden 4×5 siguiente

$$A = \begin{pmatrix} 1 & -1 & 2 & 2 & 1 \\ 3 & 3 & 0 & 2 & -1 \\ 2 & -4 & 6 & 1 & -1 \\ -1 & -3 & 2 & -3 & -2 \end{pmatrix}$$

Utilizando dicha matriz escalonada, calcular el rango de A .**Solución:**

(a) By the properties of the determinant

$$\begin{aligned} \begin{vmatrix} 1 & 0 & 2a+6 & 1 \\ 2 & -1 & 2b & 1 \\ 3 & 4 & 2c & 0 \\ 1 & -1 & 2d & 0 \end{vmatrix} &= - \begin{vmatrix} 1 & 0 & 1 & 2a+6 \\ 2 & -1 & 1 & 2b \\ 3 & 4 & 0 & 2c \\ 1 & -1 & 0 & 2d \end{vmatrix} = - \left(\begin{vmatrix} 1 & 0 & 1 & 2a \\ 2 & -1 & 1 & 2b \\ 3 & 4 & 0 & 2c \\ 1 & -1 & 0 & 2d \end{vmatrix} + \begin{vmatrix} 1 & 0 & 1 & 6 \\ 2 & -1 & 1 & 0 \\ 3 & 4 & 0 & 0 \\ 1 & -1 & 0 & 0 \end{vmatrix} \right) \\ &= -2 \begin{vmatrix} 1 & 0 & 1 & a \\ 2 & -1 & 1 & b \\ 3 & 4 & 0 & c \\ 1 & -1 & 0 & d \end{vmatrix} - (-6) \begin{vmatrix} 2 & -1 & 1 \\ 3 & 4 & 0 \\ 1 & -1 & 0 \end{vmatrix} = -20 - (-6) \times (-7) = -62. \end{aligned}$$

(b) We use Gaussian elimination to find an echelon form equivalent to the given matrix.

- row 2 $-3 \times$ (row 1), row 3 $-2 \times$ row 1, and row 4 $+$ row 1.

Exchange rows.

$$\begin{pmatrix} 1 & -1 & 2 & 2 & 1 \\ 0 & 6 & -6 & -4 & -4 \\ 0 & -2 & 2 & -3 & -3 \\ 0 & -4 & 4 & -1 & -1 \end{pmatrix} \quad \begin{pmatrix} 1 & -1 & 2 & 2 & 1 \\ 0 & -2 & 2 & -3 & -3 \\ 0 & -4 & 4 & -1 & -1 \\ 0 & 6 & -6 & -4 & -4 \end{pmatrix}$$

- Subtract row 2 multiplied by 2 to row 3, and add row 2 multiplied by 3 to row 4.

$$\begin{pmatrix} 1 & -1 & 2 & 2 & 1 \\ 0 & -2 & 2 & -3 & -3 \\ 0 & 0 & 0 & 5 & 5 \\ 0 & 0 & 0 & -13 & -13 \end{pmatrix}$$

- We arrive to the echelon form where only the bottom row is null, thus the rank of the matrix is 3.

$$\begin{pmatrix} 1 & -1 & 2 & 2 & 1 \\ 0 & -2 & 2 & -3 & -3 \\ 0 & 0 & 0 & 5 & 5 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

2

Se considera la siguiente matriz.

$$A = \begin{pmatrix} 2/3 & -1/3 \\ -1/3 & 2/3 \end{pmatrix}.$$

- (a) (10 puntos) Calcular los valores y vectores propios de A y probar que A es diagonalizable. Encontrar una de las formas diagonales D de A y la correspondiente matriz asociada P . Justificar las respuestas.
- (b) (10 puntos) Utilizando los resultados obtenidos en el apartado (a), calcular la n -ésima potencia de A , es decir, calcular

$$\begin{pmatrix} 2/3 & -1/3 \\ -1/3 & 2/3 \end{pmatrix}^n,$$

donde n es un número entero positivo. Justificar la respuesta.

Solución:

- (a) $p_A(\lambda) = |A - \lambda I| = (2/3 - \lambda)^2 - 1/9$. The roots (eigenvalues) are $\lambda = 1$ and $\lambda = 1/3$, thus A is diagonalizable since all the eigenvalues are real and distinct.

The eigenspace $S(1)$ is obtained from solving $\begin{cases} -(1/3)x - (1/3)y = 0 \\ -(1/3)x - (1/3)y = 0 \end{cases}$, that is, $S(1) = \{(x, -x) : x \in \mathbb{R}\}$, thus $(1, -1)$ is an eigenvalue associated to $\lambda = 1$.

The eigenspace $S(1/3)$ is obtained from solving $\begin{cases} (1/3)x - (1/3)y = 0 \\ -(1/3)x + (1/3)y = 0 \end{cases}$, that is, $S(1/3) = \{(x, x) : x \in \mathbb{R}\}$, thus $(1, 1)$ is an eigenvalue associated to $\lambda = 1/3$.

Hence

$$D = \begin{pmatrix} 1 & 0 \\ 0 & 1/3 \end{pmatrix}, \quad P = \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}.$$

- (b) From $D = P^{-1}AP$ we obtain $A = PDP^{-1}$ and then $A^n = PD^nP^{-1}$ as it is shown in the class notes.

The inverse of P is $P^{-1} = \begin{pmatrix} 1/2 & -1/2 \\ 1/2 & 1/2 \end{pmatrix}$.

Thus

$$\begin{pmatrix} 2/3 & -1/3 \\ -1/3 & 2/3 \end{pmatrix}^n = \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1/3 \end{pmatrix}^n \begin{pmatrix} 1/2 & -1/2 \\ 1/2 & 1/2 \end{pmatrix},$$

obtaining

$$\begin{pmatrix} 2/3 & -1/3 \\ -1/3 & 2/3 \end{pmatrix}^n = \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1/3^n \end{pmatrix} \begin{pmatrix} 1/2 & -1/2 \\ 1/2 & 1/2 \end{pmatrix}.$$

Finally

$$\begin{pmatrix} 2/3 & -1/3 \\ -1/3 & 2/3 \end{pmatrix}^n = \begin{pmatrix} 1/2 + (1/2)(1/3^n) & -1/2 + (1/2)(1/3^n) \\ -1/2 + (1/2)(1/3^n) & 1/2 + (1/2)(1/3^n) \end{pmatrix}$$

3

- (a) (10 puntos) Clasificar la forma cuadrática $Q(x, y, z) = \frac{a}{3}x^2 + ay^2 + \frac{27}{a}z^2 + axy + 3yz$, donde $a \neq 0$ es un parámetro.
- (b) (10 puntos) Representar el subconjunto del plano $D = \{(x, y) : 0 \leq x \leq 2, y \leq x, y \leq 2 - x\}$ y encontrar el valor de la integral doble siguiente:

$$\iint_D \sqrt{x+y} \, dx \, dy.$$


Solución:

- (a) The matrix associated to the quadratic form is

$$A = \begin{pmatrix} \frac{a}{3} & \frac{a}{2} & 0 \\ \frac{a}{2} & a & \frac{3}{2} \\ 0 & \frac{3}{2} & \frac{27}{a} \end{pmatrix},$$

with principal minors $A_1 = \frac{a}{3}$, $A_2 = \frac{a^2}{12}$ and $A_3 = |A| = \frac{3}{2}a$. Thus, Q is definite positive if $a > 0$ and definite negative if $a < 0$.


- (b) The set D is the infinite region represented in the figure below




To calculate the integral, note that integrand function $\sqrt{x+y}$ is defined only when $x+y \geq 0$, thus the effective region of integration is the subset of D given by

$$D' = \{(x, y) : 0 \leq x \leq 2, y \leq x, y \leq 2 - x, x + y \geq 0\}.$$


The subset D' is represented here



To integrate, we write $D' = D_1 \cup D_2$, where D_1 is the triangle of vertexes $(0, 0)$, $(1, 1)$ and $(2, 0)$ and D_2 is the triangle of vertexes $(0, 0)$, $(2, -2)$ and $(2, 0)$.



$$\begin{aligned}
 \iint_{D_1} \sqrt{x+y} \, dx \, dy &= \int_0^1 dy \int_y^{2-y} \sqrt{x+y} \, dx \\
 &= \int_0^1 \frac{2}{3} (x+y)^{3/2} \Big|_{x=y}^{x=2-y} dy \\
 &= \frac{2}{3} \int_0^1 \left(2^{3/2} - (2y)^{3/2} \right) dy \\
 &= \frac{2}{3} 2^{3/2} \int_0^1 \left(1 - y^{3/2} \right) dy \\
 &= \frac{2}{3} 2^{3/2} \left(y \Big|_{y=0}^{y=1} - \frac{2}{5} y^{5/2} \Big|_{y=0}^{y=1} \right) \\
 &= \frac{2}{3} 2^{3/2} \left(1 - \frac{2}{5} \right) \\
 &= \frac{2\sqrt{8}}{5} \left(= \frac{4\sqrt{2}}{5} \right).
 \end{aligned}$$



$$\begin{aligned}
 \iint_{D_2} \sqrt{x+y} \, dx \, dy &= \int_{-2}^0 dy \int_{-y}^2 \sqrt{x+y} \, dx \\
 &= \int_{-2}^0 \frac{2}{3} (x+y)^{3/2} \Big|_{x=-y}^{x=2} dy \\
 &= \frac{2}{3} \int_{-2}^0 \left((2+y)^{3/2} - (-y+y)^{3/2} \right) dy \\
 &= \frac{2}{3} \int_{-2}^0 (2+y)^{3/2} dy \\
 &= \frac{2}{3} \frac{2}{5} (2+y)^{5/2} \Big|_{y=-2}^{y=0} \\
 &= \frac{4}{15} \left(2^{5/2} - (2-2)^{5/2} \right) \\
 &= \frac{16}{15} \sqrt{2}.
 \end{aligned}$$

Hence

$$\iint_D \sqrt{x+y} \, dx \, dy = \iint_{D_1} \sqrt{x+y} \, dx \, dy + \iint_{D_2} \sqrt{x+y} \, dx \, dy = \frac{4}{5} \sqrt{2} + \frac{16}{15} \sqrt{2} = \frac{28}{15} \sqrt{2}.$$

4

- (a) (10 puntos) Estudiar la convergencia de la integral impropia

$$\int_3^4 \ln(x-3) dx$$

y encontrar su valor de forma razonada si resultara convergente.

Ayuda: Calcular $\lim_{x \rightarrow 3^+} (x-3) \ln(x-3)$ mediante la Regla de L'Hopital.

- (b) (10 puntos) Estudiar la convergencia de la integral impropia

$$\int_5^\infty \frac{1}{25+x^2} dx$$

y encontrar su valor de forma razonada si resultara convergente.

Solución:

- (a) We first find a primitive of the integrand. Let
- $u = \ln(x-3)$
- and
- $dv = dx$
- , so
- $du = \frac{1}{x-3} dx$
- and
- $v = x$
- . By parts,

$$\int \ln(x-3) dx = x \ln(x-3) - \int \frac{x}{x-3} dx.$$

Now, $\frac{x}{x-3} = 1 + \frac{3}{x-3}$, hence

$$\int \frac{x}{x-3} dx = \int \left(1 + \frac{3}{x-3}\right) dx = x + 3 \ln(x-3) \text{ (plus an arbitrary constant that we do not consider).}$$

Hence

$$\int \ln(x-3) dx = x \ln(x-3) - x - 3 \ln(x-3) = (x-3) \ln(x-3) - x.$$

Thus

$$\begin{aligned} \int_3^4 \ln(x-3) dx &= \lim_{a \rightarrow 3^+} \int_a^4 \ln(x-3) dx \\ &= \lim_{a \rightarrow 3^+} \left((x-3) \ln(x-3) - x \right) \Big|_{x=a}^4 \\ &= \lim_{a \rightarrow 3^+} \left((4-3) \ln(4-3) - 4 - (a-3) \ln(a-3) - a \right) \\ &= (4-3) \ln(4-3) - 4 - \lim_{a \rightarrow 3^+} \left((a-3) \ln(a-3) - a \right) \\ &= 1 \times 0 - 4 + 3 = -1, \end{aligned}$$

once we have calculated that $\lim_{x \rightarrow 3^+} (x-3) \ln(x-3) = 0$ using the hint:

$$\lim_{x \rightarrow 3^+} (x-3) \ln(x-3) = \lim_{x \rightarrow 3^+} \frac{\ln(x-3)}{\frac{1}{x-3}} = \lim_{x \rightarrow 3^+} \frac{\frac{1}{x-3}}{-\frac{1}{(x-3)^2}} = \lim_{x \rightarrow 3^+} -\frac{(x-3)^2}{(x-3)} = 0.$$

- (b) Since
- $25+x^2 \geq x^2$
- , we have
- $\frac{1}{25+x^2} \leq \frac{1}{x^2}$
- . Since
- $\int_5^\infty \frac{1}{x^2} dx$
- is convergent, we have, by the comparison principle, that the requested integral converges. Let us find its value.

$$\begin{aligned} \int_5^\infty \frac{1}{25+x^2} dx &= \frac{1}{25} \int_5^\infty \frac{1}{1+(x/5)^2} dx \\ &= \frac{1}{25} \int_1^\infty \frac{5}{1+t^2} dt \quad (t = \frac{x}{5}, dt = \frac{dx}{5}) \\ &= \frac{5}{25} \int_1^\infty \frac{1}{1+t^2} dt \\ &= \frac{1}{5} \arctan t \Big|_{t=0}^{t=1} = \frac{1}{5} (\arctan 1 - \arctan 0) = \frac{1}{5} \frac{\pi}{4} = \frac{\pi}{20}. \end{aligned}$$

5

- (a) (10 puntos) Calcular el límite siguiente:

$$\lim_{n \rightarrow \infty} \frac{\sqrt{n^2 + 7} - n}{\sqrt{n^2 + 13} - n}.$$

Justificar la respuesta.

- (b) (10 puntos) Enunciar el Teorema de Leibniz sobre la convergencia de series alternadas.

Verificar que la serie

$$\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{\sqrt[3]{n}}.$$

cumple todas las hipótesis del Teorema de Leibniz. Si S denota la suma de la serie y S_{26} denota la suma parcial de los 26 primeros sumandos de la serie infinita,

$$\frac{1}{\sqrt[3]{1}} - \frac{1}{\sqrt[3]{2}} + \cdots + \frac{1}{\sqrt[3]{25}} - \frac{1}{\sqrt[3]{26}},$$

¿cuál es la cota del error $|S - S_{26}|$ que proporciona el Teorema de Leibniz?

Solución:

- (a)

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\sqrt{n^2 + 7} - n}{\sqrt{n^2 + 13} - n} &= \lim_{n \rightarrow \infty} \frac{(\sqrt{n^2 + 7} - n)(\sqrt{n^2 + 7} + n)(\sqrt{n^2 + 13} + n)}{(\sqrt{n^2 + 13} - n)(\sqrt{n^2 + 13} + n)(\sqrt{n^2 + 7} + n)} \\ &= \lim_{n \rightarrow \infty} \frac{7(\sqrt{n^2 + 13} + n)}{13(\sqrt{n^2 + 7} + n)} \\ &= \frac{7}{13} \lim_{n \rightarrow \infty} \frac{\sqrt{1 + \frac{13}{n^2}} + 1}{\sqrt{1 + \frac{7}{n^2}} + 1} \\ &= \frac{7}{13}. \end{aligned}$$

- (b) For the statement of the Theorem of Leibniz for alternating series, see the class notes.

Let us denote $a_n = \frac{1}{\sqrt[3]{n}}$. Then $|S - S_{26}| < a_{27} = \frac{1}{\sqrt[3]{27}} = \frac{1}{3}$.