

Repeated games

2. Infinite repetition

Universidad Carlos III de Madrid

Infinite repetition

- A stage game is played at periods 1, 2, 3,... **with no end**.
- At period t players **observe** the results of all previous periods, from 1 to $t - 1$.
- Each player **discounts** future payoffs using a discount factor δ , $0 < \delta < 1$.
- The extensive form is infinite and **has no final nodes**; thus, we have to adapt the definition of a **strategy** and the **payoffs** assignment.
- A strategy is now a **rule** to assign actions to the static game in each stage of a subgame depending on the history leading to that subgame.
- Instead of assigning payoffs to final nodes, we will assign **payoffs to every possible strategy** as the discounted sum of payoffs obtained in each stage.
- Example: the **presented discounted value** of obtaining Π in each period is:


$$\Pi + \delta\Pi + \delta^2\Pi + \delta^3\Pi + \dots = \Pi \sum_{t=0}^{\infty} \delta^t = \frac{1}{1-\delta} \Pi.$$

The prisoners' dilemma

- As in the case of finite games , the number of equilibria is potentially **huge**.
- Rather than finding all of them, we will look for **interesting equilibria**, typically equilibria that sustain some payoffs.
- E.g., consider the prisoners' dilemma repeated infinitely many times: can we sustain **cooperation**?

		Player 2	
		D	C
Player 1	D	1 , 1	5 , 0
	C	0 , 5	4 , 4

Extensive form



Infinitely many repetitions

Trigger strategy

- There are **infinitely many** subgames.
- Each subgame is **identical** to the whole game.
- We are interested in sustaining **cooperation**.
- We will use **trigger** strategies:
 - Cooperate if cooperation was played at all previous stages (**prize**).
 - After a deviation, play a NE of the stage game forever (**punishment**).
- More formally, in our example:
 - At $t = 1$: Play (C, C).
 - At $t > 1$: Play (C, C) if (C, C) was played at all $t' < t$.
Play (D, D) if at some $t' < t$ (C, C) was not played.

Trigger strategy

- Let us check that the trigger strategy is indeed a SPNE.
- Two steps:
 1. Check it is a NE of the repeated game.
 2. Check it is a NE of each subgame.
- We will make use of two facts:
 - Each subgame is **identical** to the whole game.
 - The trigger strategy separates **subgames in two groups**:
 1. Subgames after a history of **cooperation**.
 2. Subgames after a history that includes **some non-cooperative** play.

Trigger strategy

At $t = 1$: Play (C, C).

At $t > 1$: Play (C, C) if (C, C) was played at all $t' < t$.

Play (D, D) if at some $t' < t$ (C, C) was not played.

- It is a NE of the **repeated game**:
- If both players **follow** the trigger strategy, each one gets:

$$4 + 4\delta + 4\delta^2 + 4\delta^3 + \dots = \frac{4}{1-\delta}$$

- If one **deviates** only in the first stage, and plays D she will get:

$$5 + \delta + \delta^2 + \delta^3 + \dots = 5 + \frac{\delta}{1-\delta}$$

- The deviation is not profitable as long as:

$$\frac{4}{1-\delta} \geq 5 + \frac{\delta}{1-\delta} \quad \text{or} \quad \delta \geq \frac{1}{4}.$$

- **Have we shown** that it is a Nash equilibrium if $\delta \geq \frac{1}{4}$?
- No: we have just shown that **one among infinitely many possible deviations** is not profitable.

Trigger strategy

At $t = 1$: Play (C, C).
 At $t > 1$: Play (C, C) if (C, C) was played at all $t' < t$.
 Play (D, D) if at some $t' < t$ (C, C) was not played.

- Which other deviations exist?:
 - Deviations in **just one period** $t > 1$.
 - Deviations in **more than one period**.
- If a player deviates to D in just one period at $t > 1$:

$$4 + 4\delta + \dots + 4\delta^{t-1} + 5\delta^t + \delta^{t+1} + \delta^{t+2} + \dots,$$

compared to non deviating:

$$4 + 4\delta + \dots + 4\delta^{t-1} + 4\delta^t + 4\delta^{t+1} + 4\delta^{t+2} + \dots,$$

If we eliminate the first t terms ($4 + 4\delta + \dots + 4\delta^{t-1}$), that are the same in both cases, and divide by δ^t , we are in the same case as when we considered deviations at $t = 1$.

- If a player deviates to D at $t = 1$ and deviates again at t_1, t_2, \dots :

$$5 + \delta + \delta^2 + \dots + \delta^{t_1-1} + 0\delta^{t_1} + \delta^{t_1+1} + \dots + \delta^{t_2-1} + 0\delta^{t_2} + \delta^{t_2+1} + \dots.$$

We observe that the payoff is smaller than deviating only at $t = 1$:

$$5 + \delta + \delta^2 + \dots + \delta^{t_1-1} + \delta^{t_1} + \delta^{t_1+1} + \dots + \delta^{t_2-1} + \delta^{t_2} + \delta^{t_2+1} + \dots.$$

- Thus, the **best deviation** is to deviate in just one period. Now we conclude that the trigger strategy is a NE of the whole game if $\delta \geq \frac{1}{4}$.

Trigger strategy

At $t = 1$: Play (C, C).
 At $t > 1$: Play (C, C) if (C, C) was played at all $t' < t$.
 Play (D, D) if at some $t' < t$ (C, C) was not played.

- Let us now show that the trigger strategy is a NE in each subgame **that is not** the whole game.
- Recall that **all subgames are equal** (the infinite repetition of the prisoners' dilemma) and that the trigger strategy separates **subgames in two groups** at the time to prescribe the actions to follow:
 - 1) Subgames after a history of **cooperation**.
 - 2) Subgames after a history that includes at least one period in which **there has been no full cooperation**.
- Subgames at t after (C, C) has been played for all $t' < t$:
 - In these subgames, the trigger strategy prescribes the **same actions** as at $t = 1$ and the subgame is the **same** as the whole game.
 - Thus, the trigger strategy is a NE in those subgames under the **same conditions** as in the whole game.
- Subgames at t if for some $t' < t$ the actions (C, C) were not played.
 - If players do not deviate, each one will get:
 $\delta^t + \delta^{t+1} + \dots + \delta^{t'-1} + \delta^{t'} + \delta^{t'+1} + \dots$
 - If one deviates at t, t', \dots :
 $0\delta^t + \delta^{t+1} + \dots + \delta^{t'-1} + 0\delta^{t'} + \delta^{t'+1} + \dots$
 - Thus, the trigger strategy is a NE in those subgames for any value of δ .
- We conclude that the trigger strategy is a SPNE of the game.

Trigger strategy

- There are many **other payoffs**, besides (1, 1) and (4, 4), that can be sustained in a SPNE. Let us see how to sustain another one.
- Consider the following **trigger strategy**:
 - At $t = \text{odd}$ play (C, C).
 - At $t = \text{even}$ play (D, C).
 - Keep playing like that if no one deviates.
 - If at some point some one deviates, play (D, D) forever after.
- If they both **follow** the strategy, Player 1 gets:

$$4 + \delta 5 + \delta^2 4 + \delta^3 5 + \dots = \frac{4}{1-\delta^2} + \frac{5\delta}{1-\delta^2}.$$

- Player 2 gets:

$$4 + \delta 0 + \delta^2 4 + \delta^3 0 + \dots = \frac{4}{1-\delta^2}.$$

- For **high enough** values of δ , the strategy is a SPNE.
- In general, we can **design** the following trigger strategy, and it will be a SPNE for high enough values of δ :
 - Define a way to play that give the **payoffs to be sustained** (they must be larger or equal to the payoffs in a NE).
 - Keep playing like that **if no one deviates**.
 - **If some one deviates**, change to play the NE forever after.

Undefined or uncertain repetition

- **Uncertain repetition**: the game continues to the next period with probability p .
- Keep the **discount** rate δ .
- The expected discount payoff of a series of payments is now calculated as:

$$\Pi_1 + p\delta\Pi_2 + (p\delta)^2\Pi_3 + (p\delta)^3\Pi_4 + \dots = \sum_{t=1}^{\infty} (p\delta)^{t-1} \Pi_t.$$

- In particular, if $\Pi_t = \Pi$ for all t :

$$\sum_{t=1}^{\infty} (p\delta)^{t-1} \Pi_t = \Pi \sum_{t=0}^{\infty} (p\delta)^t = \frac{1}{1-p\delta} \Pi.$$

- We observe that the model is equivalent to the one already studied with a discount rate $\delta' = p\delta$.

Application to Cournot oligopoly

- Firms interact infinitely many times (or many times with not a specified final):
 - They can learn no **coordinate** their strategies.
 - They can threat with some periods of **punishment** (low profits) in case of a deviation.
- Implications:
 - If firms are patient enough, they can sustain **monopoly prices** in each period.
 - The larger the number of firms, the more difficult it is to collude.

Application to Cournot oligopoly

- Recall the **Cournot** game seen in static games (2 firms, linear demand and equal, constant marginal costs:
$$\Pi_i = q_i(a - (q_1 + q_2)) - cq_i.$$
- The Nash-Cournot **equilibrium** is: $(q_1^C, q_2^C) = \left(\frac{a-c}{3}, \frac{a-c}{3}\right).$
- Equilibrium **profits** are $\Pi_i^C = \Pi_i^C = \frac{1}{9}(a - c)^2.$
- Recall also that the **monopoly quantity** maximizes profits and is $q^M = \frac{a-c}{2},$ with **profits** $\Pi^M = \frac{1}{4}(a - c)^2.$
- Thus, if each produces $q_i^M = \frac{q^M}{2} = \frac{a-c}{4},$ each one will enjoy profits equal to $\Pi_i^M = \frac{\Pi^M}{2} = \frac{1}{8}(a - c)^2,$ **higher** than profits in the Nash-Cournot equilibrium.

Application to Cournot oligopoly

- Trigger strategy:
 - Start cooperating and **keep cooperating** as long as no one deviates.
 - If someone deviates, **stop cooperating**.
- Formally:
 - At $t = 1$: Play $(q_1^M, q_2^M) = \left(\frac{a-c}{4}, \frac{a-c}{4}\right)$.
 - At $t > 1$: Play (q_1^M, q_2^M) if (q_1^M, q_2^M) was played at all $t' < t$.
 Play $(q_1^C, q_2^C) = \left(\frac{a-c}{3}, \frac{a-c}{3}\right)$ if at some period $t' < t$ (q_1^M, q_2^M) was not played.
- Differences with the repeated prisoners' dilemma:
 - There are infinitely many strategies in the stage game, not just two.
 - The best deviation in the stage game against q_j^M is not to play q_i^C , but the quantity indicated by the reaction function. Denote this best deviation by q_i^D :

$$q_i^D = \frac{a-c-q_j^M}{2} = \frac{a-c-\frac{a-c}{4}}{2} = \frac{3}{8}(a-c).$$

- Profits for Firm i after (q_i^D, q_j^M) are:

$$\Pi_i^D(q_i^D, q_j^M) = \left(a - \frac{3}{8}(a-c) - \frac{a-c}{4} - c\right) \frac{3}{8}(a-c) = \frac{9}{64}(a-c)^2.$$

Application to Cournot oligopoly

- Let us check that the trigger strategy is a SPNE:
- If both firms **follow** the trigger strategy, each one has:

$$\Pi_i^M + \delta \Pi_i^M + \delta^2 \Pi_i^M + \dots = \Pi_i^M \frac{1}{1-\delta} = \frac{1}{8}(a-c)^2 \frac{1}{1-\delta}.$$

- If Firm i **deviates only during the first period** it gets:

$$\Pi_i^D + \delta \Pi_i^C + \delta^2 \Pi_i^C + \dots = \frac{9}{64}(a-c)^2 + \frac{1}{9}(a-c)^2 \frac{\delta}{1-\delta}.$$

- The deviation is not profitable if:

$$\frac{1}{8}(a-c)^2 \frac{1}{1-\delta} \geq \frac{9}{64}(a-c)^2 + \frac{1}{9}(a-c)^2 \frac{\delta}{1-\delta}.$$

- Solving for δ we obtain that $\delta \geq \frac{9}{17}$ is the **condition for no deviations by either firm**.
- Repeating the same arguments seen in the repeated prisoners' dilemma, we can show that this is the **best deviation** and, thus, that the trigger strategy is a NE.
- Also repeating those arguments, we can show that it is also a NE in all **subgames** and, then, a SPNE.

More results with games repeated infinitely many times

- Consider again the infinitely repeated prisoners' dilemma.
- The trigger strategy is too severe and **punishes both players for too long**. We can design a strategy that limits the punishment to a **small number of periods** and that **punishes only the deviator**.
- The analysis can be extended to **more players**.
- The analysis can be extended also to the case of **casual encounters** between two random players within a society:
 - E.g.: in a society with 100 individuals, each day two players chosen at **random** play the prisoners' dilemma. All observe their behavior.
 - Rules:
 - Each player is born with the "**cooperator**" label. The label is visible to all.
 - A player keeps the label if she **cooperate with another "cooperator"** and **if does not cooperate against someone with the label "no cooperator"**.
 - If a player **does not follow** the above strategy, her label changes to "no cooperator".
 - With those rules, **cooperate with a "cooperator" and no cooperate with a "no cooperator" is a SPNE**.

Back to games repeated finitely many times

- Consider again the prisoners' dilemma repeated **finitely many** times.
- According to our analysis, the only el SPNE consists on **repeating the NEa** in each stage in all subgames.
- However, if the game is repeated, say 200 times, it is hard to think that we cannot expect **cooperation during many periods** (all except a few at the end).
- Small **deviations from the rationality assumption** allows that. These are just three examples:
 - One of the players may believe that the number of repetitions is **different** than the actual number y , and the other player may take that into account.
 - There may be **mistakes** when keeping track of the periods already played.
 - With some probability, one player may be **irrational** and always cooperate.