

Gamson's law and hedonic games

Michel Le Breton · Ignacio Ortuño-Ortín ·
Shlomo Weber

Received: 13 December 2006 / Accepted: 19 January 2007 / Published online: 4 April 2007
© Springer-Verlag 2007

Abstract This note adds one celebrated coalition formation game due to Gamson (Am Soc Rev 26:373–382 1961a, Am Soc Rev 26:565–573, 1961b) in the list of applications of the theory of hedonic games explored by Banerjee et al. (Soc Choice Welf 18:135–153, 2001) and Bogomolnaia and Jackson (Games Econ Behav 38: 204–230, 2002). We apply their results to study the original Gamson game and offer extensions both to a multi-dimensional characteristics space and to an infinite number of players.

1 Motivation

In his seminal work [Gamson \(1961a\)](#) examined a coalition formation game where each player in the “winning” coalition receives a share of the total coali-

M. L. Breton (✉)
Université de Toulouse 1, GREMAQ and IDEI,
21 allée de Brienne, 31000 Toulouse, France
e-mail: lebreton@cict.fr

I. Ortuño-Ortín
Department of Economics,
Universidad de Alicante and IVIE,
Alicante, Spain

S. Weber
Department of Economics,
Southern Methodist University,
Dallas, USA

S. Weber
CORE, Catholic University of Louvain,
Louvain La Neuve, Belgium

tion resources proportional to her own contribution. This celebrated coalition formation game, often referred as the Gamson's game or Gamson's law,¹ has generated a tremendous interest in various areas of social sciences. Gamson's research was motivated by early work of Mills (1953) and Caplow (1956) who analyzed the structure of coalitions formed in three-person families and pointed out the presence of the weakest player in the equilibrium coalition. These theoretical considerations have produced an important volume of research in *social psychology*, which made coalitional experiments² an important part of the research in this field (Bonacich et al. 1985; Chertkoff 1967, 1971; Gamson 1964). A great deal has been learned about choices of coalition partners and the way a division of coalition winnings is affected by variations in game rules and players' attributes.

As noted and illustrated by Caplow (1968), a theory of coalitions in triads could be used to analyze conflict and cooperation in many social and political environments, including *international relations*. Hsiung (1987) offers coalitional analysis of the strategic triangle of the United States, China and the Soviet Union during the Cold War era of 1950–1985. Along similar lines, Caplow (1989) discusses the failure of peace planning in 1815, 1919 and 1945, while Zagare (1984) examines the outcome of the Geneva conference on Vietnam in 1954.

Together with the minimal size principle introduced by Riker (1962), Gamson's law is also among the most popular hypothesis in *political science* and is one of the prominent landmarks in empirical models of the allocation of cabinet portfolios in coalition governments.³ Laver (1998) points out that "Gamson's law boasts one of the highest non-trivial R-squared figures in political science". One of the great appeals of Gamson's law is its intuitive nature and the parsimony it offers, being independent of the game form underlying the legislative bargaining process. The latter feature of Gamson's law continues to generate an interest in political science as evidenced by recent contributions of Carroll et al. (2004) and Fréchette et al. (2004).

The purpose of this brief note is to place the game-theoretical interpretation of Gamson's law in the context of hedonic games developed by Banerjee et al. (2001) (BKS – henceforth) (see also Bogomolnaia and Jackson 2002). In this framework the payoff of every player depends only on the composition of the coalition she belongs to. BKS show that if players share common preferences over possible coalitions, there is a stable partition of all players into disjoint coalitions. This so-called "top-coalition property" allows us to guarantee the existence and to characterize the equilibria of the Gamson's game. We then generalize the original Gamson's game and examine the case where

¹ The terminology "Gamson's law" is somewhat inaccurate as there is no stylized and universal prediction (coalitional pattern) associated with the Gamson's game. It is however kept here to maintain the link with the existing literature. We are grateful to the referee for calling our attention on that point.

² Gamson (1961b) describes the results of an experiment designed to test his theory.

³ See Brown and Franklin (1973), Brown and Frendreis (1980), Warwick and Druckman (2001), Acemoglu et al. (2006) on modified versions of Gamson's law in this context.

each player is characterized by a multiple array of characteristics. We consider two different types of a generalized Gamson's game and show that under the single division method, the BKS weak top-coalition property still applies, thus, yielding the existence of a stable coalition. However, under the double division method, a stable coalition may fail to exist. Incidentally, we offer a concept of "congruent" game and demonstrate that a game admits a stable coalition if and only if it is congruent.⁴ Finally, we consider the Gamson's game with a continuum of players and derive a necessary and sufficient condition for the existence of a stable coalition.

2 Model

Let us now introduce the Gamson's game. Consider a finite set of players $N = \{1, \dots, n\}$, where each player $i \in N$ is described by a positive parameter θ_i , interpreted as her endowment. Without loss of generality, we assume that the players are ordered according to the value of this parameter, i.e., $\theta_1 \geq \theta_2 \geq \dots \geq \theta_n$. For every $S \subset N$ denote by

$$\Theta(S) = \sum_{i \in S} \theta_i$$

the total endowment of coalition S and put $\Theta(N) = 1$.

A coalition $S \subset N$ is *winning* if the total endowment of S exceeds the total endowment of its complement $N \setminus S$. We denote by \mathcal{W} the set of winning coalitions, i.e.,

$$\mathcal{W} = \left\{ S \subset N \mid \Theta(S) > \frac{1}{2} \right\}.$$

If a winning coalition S forms, every member of S derives the payoff $U_i(S)$, which is a share of her endowment in S . A coalition which is not winning, has nothing to offer to its members. That is, for every $i \in S$

$$U_i(S) = \begin{cases} \frac{\theta_i}{\Theta(S)} & \text{if } S \in \mathcal{W} \\ 0 & \text{otherwise.} \end{cases}$$

The Gamson's game is hedonic in the sense that each player's strategic considerations take into account the payoff distribution after the coalition she belongs is formed, and thus, the payoff is the function of the composition of the formed coalition only. We now turn to the notion of stability:

⁴ We also point out that the notions of congruency and top-coalition property in general do not coincide.

Definition A partition of the set N in two coalitions S and $N \setminus S$ is called *stable* if there is no coalition T such that

$$U_i(T) > U_i(S) \quad \text{for all } i \in T \cap S \quad \text{and} \quad U_i(T) > U_i(N \setminus S) \quad \text{for all } i \in T \cap (N \setminus S).$$

Obviously, if a partition is stable, one of its two elements, say S , is a winning coalition. For the above inequality to hold true, it must be the case that T is winning. Therefore, we deduce that $T \cap S \neq \emptyset$ and that the second part part of the test can be deleted. Thus, a partition is stable if and only if there is no winning coalition T with $U_i(T) > U_i(S)$ for all $i \in T \cap S$. We have the following observation:

Result 2.1 A winning coalition S is stable if and only if it is weakest (in terms of endowment) among all winning coalitions:

$$\Theta(S) = \min_{T \in \mathcal{W}} \Theta(T).$$

A simple direct proof of this result can be provided. However, it immediately follows from Corollary 2 in BKS.

It is worth pointing out that the hedonic framework of BKS allows for an extension of the Gamson's game. Consider a positive-valued function $H : \mathfrak{R}_+ \times \mathfrak{R}_+ \times 2^N \rightarrow \mathfrak{R}_{++}$, which is increasing in the first argument and is decreasing in the second. Put, for all $i \in S$

$$U_i(S) = \begin{cases} H(\theta_i, \Theta(S), S) & \text{if } S \in \mathcal{W} \\ 0 & \text{otherwise} \end{cases}$$

Since this generalized game satisfies the *weak top-coalition property*, the main theorem in BKS yields the existence of a stable coalition.⁵ In the case of a triad, an equilibrium coalition contains the weakest player:

Result 2.2 Let $n = 3$ and $\frac{1}{2} > \theta_1 > \theta_2 > \theta_3$, i.e., no singleton is a winning coalition. Then, the unique stable coalition is $\{2, 3\}$.

However, if $n \geq 4$, the stable coalition does not necessarily contain the weakest player. Consider the following example:

Example 2.3 Let $n = 4$ and $\theta_1 = 0.40$, $\theta_2 = 0.27$, $\theta_3 = 0.25$ and $\theta_4 = 0.08$. Then the unique stable coalition is $\{2, 3\}$, which does not contain player 4.

Finally, one can raise the question whether stable coalitions are *connected* or *consecutive* (Greenberg and Weber 1986), where, to recall, a coalition S is connected if for every three different players, $i < j < k$, $i, k \in S$ implies that j

⁵ Note, however, that this formulation allows for increasing returns to scale in terms of coalition size. Thus, a stable winning coalition is not necessarily minimal. Note that this framework covers the case where coalitions are not necessarily either winning or losing. In the case where a surplus is also distributed among small coalitions, a core stable partition may contain more than two elements.

belongs to S as well. Assume now that the game is *strict*, i.e., there is no coalition S with $\Theta(S) = \frac{1}{2}$. We have

Result 2.4 Let $n = 4$ and $\frac{1}{2} > \theta_1 > \theta_2 > \theta_3 > \theta_4$. If a stable coalition S does not contain the weakest player, it is connected.

Proof Suppose there is a stable coalition S that does not contain player 4 and is not connected. Then $S = \{1, 3\}$. Since S is stable and the game is strict, $\theta_2 + \theta_3 < \frac{1}{2}$. But this implies that $\theta_1 + \theta_4 > \frac{1}{2}$ and $\{1, 4\}$ is a stable coalition, a contradiction. □

However, Result 2.4 cannot be extended to a larger number of players:

Example 2.5 Let $n = 5$ and $\theta_1 = 0.40, \theta_2 = 0.30, \theta_3 = 0.12, \theta_4 = 0.11$ and $\theta_5 = 0.07$. Then the unique stable coalition is $\{1, 4\}$.

We leave open the interesting problem of the characterization of vectors $\theta \equiv (\theta_1, \theta_2, \dots, \theta_n)$ for which stable coalitions exhibit connectedness or include the weakest player.

In the next section we examine a multi-dimensional variant of the Gamson’s game, where each player is characterized by a multiple array of characteristics.

3 Multi-characteristic extension

Note that Gamson’s law has been formulated for the weighted majority games and we could conceivably consider an extension to the case where, unlike in the previous section, each player $i \in N$ is identified by $K > 1$ characteristics, specifically, by the vector $\theta_i = (\theta_i^1, \theta_i^2, \dots, \theta_i^K) \in \mathfrak{R}_+^K$. A winning coalition S must satisfy

$$\Theta^k(S) = \sum_{i \in S} \theta_i^k > \frac{\Theta^k(N)}{2} \quad \text{for every } k = 1, 2, \dots, K.$$

This extension of weighted majority games is relevant to describe various voting environments. An important illustration is provided by the qualified majority provisions in the treaty of Nice for which $K = 3$ and every country i is described by the vector $\Theta_i = (1, \theta_i^2, \theta_i^3)$, where θ_i^2 is the country i ’s population, and θ_i^3 is an assigned voting weight. Thus, in forming a winning coalition one takes into account the number of countries it contains as well as their total population and aggregated voting weight (Felsenthal and Machover 2001). The more recent qualified majority decision rules for the Council of Ministers of the EU that were included in the draft European Constitution proposed by the 2003 European Convention reduced the number of parameters to $K = 2$ (Felsenthal and Machover 2004). Another example is provided by the amendment of Canada’s Constitution Act of 1982 (Kilgour 1983).

There are several ways to extend the division of surplus in the Gamson's game. Let $\alpha = (\alpha^1, \alpha^2, \dots, \alpha^K)$, be the vector of positive weights assigned to each of K characteristics. Assume that $\sum_{k=1}^{k=K} \alpha^k = 1$.

First, consider a *single division method*, where the payoff $U_i(S)$ of every member i of the winning coalition S is the share of her "weighted" endowment in S . As before, a coalition which is not winning, has nothing to offer to its members. Thus, for every $i \in S$ we have

$$U_i(S) = \begin{cases} \frac{\sum_{k=1}^{k=K} \alpha^k \theta_i^k}{\sum_{k=1}^{k=K} \alpha^k \Theta^k(S)} & \text{if } S \in \mathcal{W} \\ 0 & \text{otherwise.} \end{cases}$$

The extension of Result 2.1 to this setting is straightforward.

Result 3.1 Under the single division method, a winning coalition S is stable if it has the smallest total endowment among all winning coalitions:

$$\Theta(S) = \min_{T \in \mathcal{W}} \sum_{k=1}^{k=K} \alpha^k \Theta^k(S).$$

The situation is quite different if instead of weighting the different characteristics, the pie is divided according to the *double division* method. Namely, for each characteristic k the share of each player is determined according to the unidimensional Gamson rule. The total share is then calculated as a weighted average by utilizing the weights α^k for each characteristic k . Thus, if a winning coalition S forms, every member i of S derives the payoff $V_i(S)$, defined as follows:

$$V_i(S) = \begin{cases} \sum_{k=1}^{k=K} \alpha^k \frac{\theta_i^k}{\Theta^k(S)} & \text{if } S \in \mathcal{W} \\ 0 & \text{otherwise.} \end{cases}$$

Interestingly enough, this game does not always admit an equilibrium.

Result 3.2 Under the double division method, a stable coalition may fail to exist.

Proof Consider the following example with three players and three characteristics. Let $\theta_1 = (0.40, 0.35, 0.25)$, $\theta_2 = (0.25, 0.40, 0.35)$, $\theta_3 = (0.35, 0.25, 0.40)$ and $\alpha = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$. The winning coalitions consists of all groups with at least

two players. Straightforward calculations lead to:

$$\begin{aligned}
 V_1(\{1, 2\}) &= \frac{1}{3} \left(\frac{40}{75} + \frac{25}{65} + \frac{35}{60} \right) \simeq 0.50043 > 0.49957 \simeq V_1(\{1, 3\}) \\
 &= \frac{1}{3} \left(\frac{40}{65} + \frac{25}{60} + \frac{35}{75} \right) \\
 V_2(\{2, 3\}) &= \frac{1}{3} \left(\frac{40}{75} + \frac{25}{65} + \frac{35}{60} \right) \simeq 0.50043 > 0.49957 \simeq V_2(\{1, 2\}) \\
 &= \frac{1}{3} \left(\frac{40}{65} + \frac{25}{60} + \frac{35}{75} \right) \\
 V_3(\{1, 3\}) &= \frac{1}{3} \left(\frac{40}{75} + \frac{25}{65} + \frac{35}{60} \right) \simeq 0.50043 > 0.49957 \simeq V_3(\{2, 3\}) \\
 &= \frac{1}{3} \left(\frac{40}{65} + \frac{25}{60} + \frac{35}{75} \right)
 \end{aligned}$$

A stable coalition does not exist due to the emergence of the Condorcet cycle among winning coalitions. □

The last example suggests that the existence of stable coalitions calls for some degree of congruence among a subset of players. The following notion of congruence which applies to any game where the economies of scale are fully described by a proper⁶ set of winning coalitions \mathcal{W} (i.e. at equilibrium, if any, a single coalition forms) slightly generalizes the notion of weak-top coalition property introduced by BKS. In fact, congruence is necessary and sufficient for stability.

Definition A coalition $S \subseteq N$ is *congruent* if for all $i \in S$ and all $T \subseteq N$ the inequality $U_i(T) > U_i(S)$ implies that there exists $j \in T \cap S$ such that $U_j(S) \geq U_j(T)$. A coalition formation game is called *congruent* if there exists a congruent coalition.

Since we consider proper simple games, there is no loss of generality in considering partitions consisting of two coalitions,⁷ S and $N \setminus S$, where one, say S , is winning. Like for the Gamson’s game, we will refer to such coalition as a stable coalition. We have the following result:

Result 3.3 Let \mathcal{W} be a proper family of winning coalitions. A coalition formation game is stable (i.e., admits a stable coalition) if and only if it is congruent.

Proof Let the game be congruent. We show that the partition of the set N into coalitions S and $N \setminus S$, where S is congruent, is stable. Indeed, if not, then there is

⁶ A simple game (N, \mathcal{W}) is proper if $S, T \in \mathcal{W}$ implies $S \cap T \neq \emptyset$.

⁷ If the simple game is not proper, then a stable coalition structure may consist of more than two coalitions. A standard example of such assertion is provided by hedonic matching games when the population of at least six players is divided into two types.

a winning coalition T such that $U_i(T) > U_i(S)$ for all $i \in T \cap S$, a contradiction to the congruency of S .

Now, let S be a stable coalition. We show that S is congruent. Indeed, let $i \in S$ and $T \subseteq N$ be such that $U_i(T) > U_i(S)$. Since $U_j(T) > U_j(N \setminus S) = 0$ for all $j \in T \cap (N \setminus S)$, we deduce that $U_j(S) \geq U_j(T)$ for some $j \in T \cap S$. Otherwise, we would contradict our assumption that S is stable. \square

The notion of congruence generalizes the notion of weak top-coalition property in BKS as it does not imply that a congruent coalition S would be ranked at the top by all members of S . Consider the following example:

Result 3.4 Let $n = 4$ and

$$\begin{aligned} V_1(\{1, 2, 4\}) &> V_1(\{1, 2, 3\}) > V_1(T) \text{ for every other } T \text{ that includes } 1, \\ V_2(\{2, 3, 4\}) &> V_2(\{1, 2, 3\}) > V_2(T) \text{ for every other } T \text{ that includes } 2, \\ V_3(\{1, 3, 4\}) &> V_3(\{1, 2, 3\}) > V_3(T) \text{ for every other } T \text{ that includes } 3, \\ V_4(\{1, 2, 4\}) &> V_4(\{1, 2, 3\}) > V_4(T) \text{ for every other } T \text{ that includes } 4. \end{aligned}$$

Let \mathcal{W} be the set of majority coalitions. It is easy to see that $S = \{1, 2, 3\}$ is a congruent coalition. However, S does not satisfy the weak top-coalition property, as it represents only the second best choice for the members of S .

In the next section we discuss an extension of the Gamson's game to environments that consist of infinite number of players.

4 Atomless environments

The analysis in previous sections has been conducted under the assumption that the set of players was finite. Consider instead an environment with a continuum of players given by the unit interval $[0, 1]$ and a vector $\theta = (\theta^1, \theta^2, \dots, \theta^K)$ of K positive random variables, where $\theta(t) = (\theta^1(t), \theta^2(t), \dots, \theta^K(t))$ denotes the characteristics of a player of type $t \in [0, 1]$. Without loss of generality we assume that $\int_0^1 \theta^k(t) dt = 1$ for all $k = 1, \dots, K$. Given a small positive number ε ,⁸ consider a proper simple game \mathcal{W}_ε defined as follows:

$$S \in \mathcal{W}_\varepsilon \quad \text{if and only if } \theta^k(S) = \int_S \theta^k(t) dt \geq \frac{1}{2} + \varepsilon \quad \text{for all } k = 1, \dots, K.$$

We assume that the fraction $\frac{1}{K}$ of the pie is divided among the members of the coalition on the basis of their k^{th} characteristic by using the double division

⁸ The existence of a stable coalition is not guaranteed in the case of the classical majority simple game, as the game is discontinuous at $\varepsilon = 0$.

method. That is, if a coalition S forms, every player of type t of S derives the payoff $V_t(S)$, defined as follows:

$$V_t(S) = \begin{cases} \frac{1}{K} \sum_{k=1}^{k=K} \frac{\theta^k(t)}{\theta^k(S)} & \text{if } S \in \mathcal{W}_\varepsilon \\ 0 & \text{otherwise.} \end{cases}$$

An obvious sufficient condition for a coalition S to be stable is

$$\theta^k(S) = \frac{1}{2} + \varepsilon \tag{1}$$

for all $k = 1, \dots, K$.

In the finite case, condition (1) is not necessary for stability as illustrated by the following example:

Example 4.1 Let $n = 4, K = 2, \theta^1 = (0.40, 0.32, 0.17, 0.11)$ and $\theta^2 = (0.17, 0.10, 0.31, 0.42)$. Then, $\{1, 4\}$ is the unique stable coalition. However, $\theta^2(\{1, 4\}) = 0.59 > \theta^2(\{2, 4\}) = 0.52$.

The situation is however different when the set of players is atomless:

Result 4.2 There exists a stable coalition. Moreover, S is stable if and only if it satisfies (1).

Proof The fact that condition (1) is necessary and sufficient condition for stability is immediate. For the existence of a stable coalition, consider the K -dimensional valued measure μ defined as follows:

$$\mu(S) = \begin{pmatrix} \theta^1(S) \\ \theta^2(S) \\ \dots \\ \dots \\ \theta^K(S) \end{pmatrix}$$

Since μ is atomless, $\mu(\emptyset) = (0, 0, \dots, 0)$ and $\mu([0, 1]) = (1, 1, \dots, 1)$. Thus, by the Lyapunov’s theorem⁹ that there is $S \subseteq [0, 1]$ such that

$$\mu(S) = \left(\frac{1}{2} + \varepsilon, \frac{1}{2} + \varepsilon, \dots, \frac{1}{2} + \varepsilon \right).$$

□

Result 4.2 indicates that the existence of an equilibrium coalition is a relatively simple question in the case where we have a continuum of players. In fact,

⁹ This theorem asserts that the range of an atomless vector valued measure is a convex set (see, for instance, Halmos (1950) for a formal statement).

the argument above applies to any hedonic game where the payoff of a player of type $t \in [0, 1]$ is defined by

$$V_t(S) = \begin{cases} U(\theta^1(S), \theta^2(S), \dots, \theta^K(S)) & \text{if } S \in \mathcal{W} \\ 0 & \text{otherwise,} \end{cases}$$

where U is a decreasing function and there exist values $\gamma^k \in \left(\frac{1}{2}, 1\right]$, $k = 1, \dots, K$ such that the set of winning coalitions \mathcal{W} is determined by

$$\mathcal{W} = \{S \subset [0, 1] \mid \theta^k(S) \geq \gamma^k \text{ for all } k = 1, \dots, K\}.$$

5 Concluding remarks

We conclude this note by suggesting yet another extension of the Gamson's game which could accommodate a wider spectrum of applications to coalitional politics. In the model considered in this note, the outcome of the game is determined by the division of a surplus among a subset of players. As we already pointed out, in politics it may represent a cabinet portfolio allocation in coalition governments. However, governments must also choose their policy. Assuming that the ideological space is represented by a multidimensional Euclidean space and that each player i , is identified not only by her weight θ^i but also by her ideology, it would be of interest to examine hedonic games where in addition to a division of the surplus, a winning coalition is described by a specific policy depending upon the weights and ideologies of its members.¹⁰ It seems to offer a promising avenue for further research.

Acknowledgements We thank an anonymous referee for insightful comments and suggestions on the previous version of this note.

References

- Acemoglu D, Egorov G, Sonin K (2006) Coalition formation in political games. mimeo
 Banerjee S, Konishi H, Sömnez T (2001) Core in a simple coalition formation game. *Soc Choice Welf* 18:135–153
 Bogomolnaia A, Jackson MO (2002) The stability of hedonic coalition structures. *Games Econ Behav* 38:201–230
 Bonacich P, Grusky O, Peyrot M (1985) Family coalitions: a new approach and method. *Soc Psychol Q* 44:42–50
 Brown EC, Franklin MN (1973) Aspects of coalition payoffs in european parliamentary democracies. *Am Polit Sci Rev* 67:453–469
 Brown EC, Frensdreis JP (1980) Allocating coalition payoffs by conventional norm: an assessment of the evidence from cabinet coalition situations. *Am J Polit Sci* 24:753–768
 Caplow T (1956) A theory of coalitions in the triad. *Am Sociol Rev* 21:489–493
 Caplow T (1968) Two against one: coalitions in triads. Prentice-Hall, Englewood Cliffs
 Caplow T (1989) Peace games. Wesley University Press, Middletown

¹⁰ This is the setting considered by Jackson and Moselle (2002).

- Carroll R, Cox GW, Pochon M (2004) Gamson's law: how governments allocate offices. Mimeo, UC San Diego
- Chertkoff JM (1967) A revision of caplow's coalition theory. *J Exp Soc Psychol* 3:172–177
- Chertkoff JM (1971) Sociopsychological theories and research on coalition formation. In: Groenings S, Kelley EW, Leiserson M (eds) *The study of coalition behavior*. Holt, Rinehart and Winston, New York
- Felsenthal DS, Machover M (2001) The treaty of nice and qualified majority voting. *Soc Choice Welf* 18:431–464
- Felsenthal DS, Machover M (2004) Analysis of QM rules in the draft constitution for Europe proposed by the European Convention, 2003. *Soc Choice Welf* 23:1–20
- Fréchette GR, Kagel JH, Morelli M (2004) Gamson's Law versus non-cooperative bargaining theory. Mimeo, Ohio State University
- Gamson WA (1961a) A theory of coalition formation. *Am Sociol Rev* 26:373–382
- Gamson WA (1961b) An experimental test of a theory of coalition formation. *Am Sociol Rev* 26:565–573
- Gamson WA (1964) Experimental studies of coalition formation. In: Berkowitz L (ed) *Advances in experimental social psychology* Academic, New York
- Greenberg J, Weber S (1986) Strong Tiebout equilibrium under restricted preferences Domain. *J Econ Theory* 38:101–117
- Halmos P (1950) *Measure theory*. Van Nostrand, Princeton
- Hsiung JC (1987) Internal dynamics in the sino-Soviet-US Triad. In: Kim IJ (ed) *The Strategic Triangle*. Paragon, New York
- Jackson MO, Moselle B (2002) Coalition and party formation in a legislative voting game. *J Econ Theory* 103:49–87
- Kilgour DM (1983) A formal analysis of the amending formula of Canada's constitution act, 1982. *Can J Polit Sci* 16:771–777
- Laver M (1998) Models of government formation. *Ann Rev Polit Sci* 1:1–25
- Mills TM (1953) Coalition patterns in three person groups. *Am Sociol Rev* 19:657–667
- Riker WH (1962) *The theory of political coalitions*. Yale University Press, New Haven
- Warwick PV, Druckman JN (2001) Portfolio salience and the proportionality of payoffs in coalition governments. *Br J Polit Sci* 31:627–649
- Zagare FC (1984) *Game theory: concepts and applications*. Sage publications, Beverly Hills